

1 Public knowledge of alien species: a case study on aquatic biodiversity in North Iberian
2 rivers

3
4 Laura Clusa*¹, Laura Miralles¹, Sara Fernández¹, Eva García-Vázquez¹, Eduardo
5 Dopico²

6
7 1- Department of Functional Biology, University of Oviedo.

8 2- Department of Education Sciences, University of Oviedo.

9
10 **ABSTRACT**

11
12 Biological invasions have increased in recent decades due to globalization and human
13 activities. These invasions are currently one of the main threats to biodiversity, and
14 their early detection is essential for a rapid and effective response. Here, we explored
15 the use of citizen science strategies to create an early alert to detect invasive species.
16 Our main objective was to evaluate the general knowledge of volunteer participants of
17 invasive freshwater species in Asturias (north of the Iberian Peninsula) and compare it
18 with both real data from electrofishing surveys and official data from the regional
19 government. A total of 140 volunteer surveys were conducted in four different rivers
20 in Asturias. The largest group of participants consisted of males older than 50 years.
21 Four species were identified as native to the four rivers: *Anguilla anguilla*; *Mugil*
22 *cephalus*; *Salmo salar*; and, *Salmo trutta*. More than 50% of the native species
23 surveyed by electrofishing were recognized by the locals in each river region. A total
24 of 22.86% of the volunteers were able to correctly name an exotic species, and a total
25 of 7 were correctly identified: *Procambarus clarkii*; *Trachemys scripta*; *Cyprinus*
26 *carpio*; *Esox lucius*; *Salvelinus fontinalis*; *Carassius auratus*; and, *Oncorhynchus*
27 *mykiss*. However, compared to the list of actual exotic species surveyed, less than
28 40% were recognized in the four rivers. Despite the poor correlation between local
29 knowledge and real exotic aquatic fauna, citizens were able to detect one exotic
30 species not yet found in the wild in this region (*T. scripta*). Finally, more than 70% of
31 the volunteers were in favor of fighting against invasive species, although only
32 22.86% were able to identify any specific exotic species found in the region. The
33 positive attitude to exotic species control was correlated with both the level of native
34 species knowledge and the concern about the ability of exotic species to impact native
35 fauna in the region. Better training will improve public awareness, reduce the non-
36 intentional release of non-native species, and increase the detection of non-indigenous
37 species. The attitudes of the citizens make the region a promising candidate for
38 education efforts to reduce alien species introductions and help preserve fauna
39 biodiversity.

40
41 *Corresponding author: Laura Clusa, lauraclusa@gmail.com, orcid.org/0000-0002-
42 3503-2331.

43
44 Keywords: citizen science, Non indigenous species, early detection, biodiversity

45
46 The final version of the manuscript is available in:
47 <https://doi.org/10.1016/j.jnc.2018.01.001>

48
49 © <2018>. This manuscript version is made available under the CC-BY-NC-ND 4.0
50 license <http://creativecommons.org/licenses/by-nc-nd/4.0/>

51 1. Introduction

52
53 In the last several decades, the number of non-indigenous species (NIS) has increased
54 due to globalization and human activities (Hulme, 2009). NIS (exotic, alien, non-native
55 species) are species outside their native range that are often introduced by human
56 activities (introduced species). In some cases, these NIS can proliferate, undergo
57 exponential population increases and spread quickly to become invasive species
58 (Occhipinti-Ambrogi & Galil, 2004). Invasive species cause numerous impacts to the
59 community structure and environment (Chown et al., 2015). Eradication of these exotic
60 species may allow for the recovery of the native fauna, but if they become invasive,
61 eradication is nearly impossible; therefore, the main efforts should be performed before
62 establishment. Early detection is essential for rapid response and prevention from
63 further spread (Havel, Kovalenko, Thomaz, Amalfitano, & Kats, 2015).

64 During the last twenty years, the number of research articles based on citizen
65 science has increased exponentially. There are many projects involving citizen science,
66 including ones about climate change, conservation biology, ecological restoration, water
67 quality, invasive species and many more topics (Silvertown, 2009). Technology has
68 facilitated citizen science programs through new smartphone applications or on the
69 Internet. Such applications can improve communication between scientists and citizens,
70 as daily citizen observations can be easily uploaded online and made accessible to
71 researchers, thereby generating thousands of data records (Newman et al., 2012). In
72 citizen science programs, many volunteers can cover large regions with their
73 observations and help identify migration patterns, the spread of infectious diseases and
74 other ecological phenomena at a large scale (Devictor, Whittaker, & Beltrame, 2010;
75 Dickinson, Zuckerberg, & Bonter, 2010). For example, many citizen science programs
76 are the basis of large bird inventories (Tulloch, Possingham, Joseph, Szabo, & Martin,
77 2013), such as the eBird program in North America that collects five million bird
78 observations per month (Sullivan et al., 2014). Another example is the study of hosts'
79 natural resistance to virulent forest pests, which was carried out in the USA with the
80 help of citizen science (Ingwell & Preisser, 2011). Citizen science programs have been
81 increasingly used to collect data for monitoring invasive species in real time (Crall
82 et al., 2010; Dickinson et al., 2012), especially for early alerts of new NIS. Citizens can
83 reach locations that may not be accessible to scientists; for example, some areas in
84 South Florida where Burmese pythons (*Python molurus bivittatus*) have been found
85 (Falk, Snow, & Reed, 2016). Mapping crab invasions (*Carcinus maenas* and
86 *Hemigrapsus sanguineus*) along US coasts would not be possible without citizen
87 science (Delaney, Sperling, Adams, & Leung, 2008) due to the large size and extent of
88 the invaded spaces.

89 Many of the established citizen science programs are aimed at invasive plants.
90 After training, citizen scientists are able to distinguish invasive plants and collect robust
91 data, as has been shown in Texas (Gallo & Waitt, 2011). Other examples include the
92 monitoring of invasive plants in a natural reserve in Georgia, which was carried out by
93 citizens with the help of smartphones and a geo-referencing application (Hawthorne et
94 al., 2015); the Invasive Plant Atlas of New England (IPANE) that was created in 2001
95 (Bois, Silander, & Mehrhoff, 2011; Crall et al., 2011); the Invasive Plant Atlas of the
96 Mid-South (IPAMS); and the Cactus Moth Detection and Monitoring Network
97 (CMDMN) (Simpson et al., 2009). Plants are not the only species studied; successful
98 programs are also running for the detection of invasive animals, as reported above for
99 serpents and crabs. In Japan, 300,000 bumblebees were removed from the wild within
100 the monitoring program of the invasive *Bombus terrestris* (Kobori et al., 2016). In

101 North Italy and Switzerland, it was possible to map the stink bug *Halymorpha halys* and
102 develop identification guides to help track this invasive species in other regions
103 (Maistrello, Dioli, Bariselli, Mazzoli, & Giacalone-Forini, 2016). The use of citizen
104 science to detect invasive aquatic species has also increased in recent years. In Alaska,
105 citizen science was employed to control marine invasions, which are a threat to native
106 marine resources (<https://seagrant.uaf.edu/research/projects/summary.php?id=939>). In
107 Greece, 86 observations of 28 alien species reported in 2012 demonstrated the spread of
108 more than 20 invasive species (Zenetos, Koutsogiannopoulos, Ovalis, & Poursanidis,
109 2013). Citizen scientists contributed to the detection of the invasive lionfish in the
110 Caribbean Sea (Carballo-Cárdenas & Tobi, 2016). The first report of the sergeant major
111 (*Abudefduf saxatilis*) in the Mediterranean Sea was collected through citizen science on
112 the "Seawatchers" webpage (<http://www.observadoresdelmar.es>), where volunteers
113 collect data and inform scientists about new invasive species (Azzurro, Broglio,
114 Maynou, & Bariche, 2013).

115 Evaluating public knowledge of the specific taxa to be monitored is
116 recommended before creating a citizen science program about NIS. García-Llorente,
117 Martín-López, González, Alcorlo and Montes (2008) studied how different groups
118 (tourists, conservation professionals, local users and others) perceived the impact caused
119 by IAS (invasive alien species) in the Natural Reserve of Doñana (Southwest Spain) and
120 their attitudes towards IAS eradication. As many as 97% of the people in all groups
121 agreed that IAS eradication was necessary, but they were principally concerned with the
122 recent invasions and the species that had been objects of particular campaigns and
123 appeared in the news. The authors concluded that the general knowledge of citizens is
124 crucial to generating public demand for actions against invasive species, and they
125 emphasized the low concordance found between official data, real data and citizen
126 perceptions.

127 The main objective of this study was to evaluate the public's knowledge about
128 freshwater NIS in Asturias (north of the Iberian Peninsula) through a survey on species
129 reports, and the survey results were compared with actual local fauna and official data
130 from the regional and national environmental authorities. The results served to identify
131 knowledge gaps that could be used to focus training efforts in future citizen science
132 programs on aquatic biodiversity inventories.

134 2. Materials and Methods

136 2.1. Sampling sites and river biota

137
138 During 2016, four different rivers in Asturias (south-central Bay of Biscay) were
139 selected for social and biodiversity surveys: Raíces; Piles; Negro; and, Nalón (Figure 1).
140 Three of the rivers are short coastal streams (the Negro, Piles and Raíces rivers are 20,
141 16 and 15 km in length, respectively), and the Nalón River (140 km long) originates
142 from the Nalón-Narcea basin, which is the largest freshwater system in the region.
143 Sampling sites were set within river towns at the following coordinates: Luarca
144 (43.544240N, 6.535308W) on the Negro River; Salinas (43.566852N, 5.962669W) on
145 the Raíces River; Gijón (43.537846N, 5.639280W) on the Piles River; Las Caldas
146 (43.330988N, 5.930960W) and, Rioseco (43.218977N, 5.454763W) on the Nalón
147 River.

148 The most recent official inventory of the native fauna and NIS of the regional
149 rivers was published by De la Hoz (2006).

150

2.2. Social survey

A total of 140 local participants were interviewed across the study region, including males and females older than 20 years. The samples represented 0.05% of the population inhabiting the study areas. Potential interviewees were approached along recreational promenades near the rivers, and eligible and willing participants were interviewed in Spanish, their native language. Interview sessions were no longer than 5 minutes per person to facilitate easy and spontaneous responses.

The questionnaire was inspired by García-Llorente et al. (2008). The interview was formulated as a conversation to help the volunteers feel more comfortable and answer without any pressure. The survey was divided into two sections (Supplementary file 1), as follows:

- 1) General knowledge of aquatic species in the region, where the volunteers listed the species they remembered from the local river by their common names and classified them as native or exotic using their knowledge. The translation from common name to scientific name for each species was performed by the researchers. There was no possibility of error since the common names are unique for each species in this region, and there are no local variants in different valleys;
- 2) Awareness and concerns about exotic species, which contained four questions (Supplementary file 1). A final open question about the perceived changes in the river ecosystem, if any, was posed.

Pictures of animals inhabiting Iberian rivers were available if needed for recognition of a species but were not offered beforehand. The survey was previously tested in a pilot sample (N=10) to refine the questions and ensure the content was clear and easy to understand.

The word "invasive" was avoided in the interview because it has a negative connotation, so the answers from the participants were not influenced. If needed to clarify a participant's understanding, exotic species were defined as "species that are not native to this place".

The participant's answers were recorded in writing. After finishing the interview, the participants were asked to check their answers and confirm they were correctly recorded.

2.3. Ethics statement

All volunteers agreed to participate in the study and signed the informed consent for the use of their answers in research. The study was approved by the Ethics Committee from the Principality of Asturias with the permit of reference number 99/16.

2.4. Electrofishing surveys

The actual local aquatic fauna occurring in the four rivers considered was surveyed in March 2017. The standard protocol approved by the Spanish Ministry of Agriculture, Fisheries and Environment for implementing the EU Water Framework Directive 2000/60/CE was employed. This protocol, ML-R-FI-2015 (NIPO: 280-15-122-6), is based on electrofishing. The survey was carried out by Taxus S.L., a company authorized for aquatic biodiversity surveys in the Principality of Asturias. Due to the

200 different river sizes, electrofishing was carried out from one sampling site in each of the
201 three small rivers and six sampling sites along the Nalón River.

202

203 **2.5. Data analysis**

204

205 Participants were grouped by river (Nalón, Negro, Piles and Raíces), age (older or
206 younger than 50 years) and gender. Some participants provided additional information
207 about terrestrial species, but answers about only aquatic species were considered.

208 Knowledge was measured as the concordance between a participant's answer
209 and the official list of native and exotic species in Asturias, which is available in De la
210 Hoz (2006). Four measurements were obtained: the number of native species correctly
211 identified (correct natives, C_N); the number of exotic species correctly identified
212 (correct exotics, C_E); the number of exotic or absent species mistaken as native
213 (incorrect natives, I_N); and, the number of native or absent species mistaken as exotics
214 (incorrect exotics, I_E).

215 A knowledge index (K_i) was calculated as the mathematically averaged
216 knowledge (scored as correct – incorrect species) of native and exotic species, using the
217 following formula:

218

$$219 K_i = \frac{(C_N - I_N) + (C_E - I_E)}{2}$$

220

221 In the second section of the survey, the scores were 0 (I don't know), 1 (No), and
222 3 (Yes) for the questions with three answer choices; and 0 (I don't know), 1 (No), 2
223 (Sometimes, depending on the species), and 3 (Yes) for the questions with four answer
224 choices. In question C (changes in the ecosystem), the answers were classified into four
225 large groups: "Water quality", including changes in water quality (cleaner or more
226 polluted water, more or less algae, increase of floods, more sediments, lower water
227 flow...); "Fauna", including changes in aquatic fauna (for example, reduced trout
228 spawning, changes in species abundance such as an increase of *Mugil cephalus* and a
229 decrease of *Salmo trutta*); "Infrastructure", including new ponds, dams and
230 promenades; "Environment", including cleaner or dirtier surrounding environment and
231 changes in riverbank vegetation and excluding changes in the water considered in the
232 above category.

233

234 **2.6. Statistical analysis**

235

236 The data were analyzed with the program Past 3.15 (Hammer, Harper, & Ryan, 2001).
237 Normality was checked using Shapiro-Wilk W and Anderson-Darling A tests.
238 Comparisons among groups (rivers, ages or gender) were conducted using ANOVA or
239 Kruskal-Wallis to test for differences among the means or medians of the groups,
240 respectively (the latter in case of significant deviation from normality). Pairwise
241 correlations (between questions, or between knowledge and perception/opinion) were
242 calculated using Spearman's r_s . Statistical significance was set at $p < 0.05$. Bonferroni
243 correction of the significance level was applied for multiple comparisons.

244

245 **3. Results**

246

247 In total, 58 women (41.43%) and 82 men (58.57%) participated in the survey (Table 1).
248 The largest age group (34.29% of the total sample) was older than 60 years.

249 The native species identified by participants were the European freshwater
250 crayfish (“cangrejo de río” in Spanish) *Austropotamobius pallipes*, the European eel
251 (“anguila”) *Anguilla anguilla*, the sea lamprey (“lamprea”) *Petromyzon marinus*, the
252 European sea bass *Dicentrarchus labrax* (“lubina”), the flathead gray mullet *Mugil*
253 *cephalus* (“muil”, an Asturias linguistic variant, or “mújol” in standard Spanish), the
254 Atlantic salmon *Salmo salar* (“salmón”), the brown trout *Salmo trutta* (“trucha”) and
255 the gilthead sea bream *Sparus aurata* (“dorada”) (Supplementary file 2). More than 23
256 participants in each river recognized *S. trutta* as a native species. Several exotic species
257 introduced from other Spanish regions were considered to be native by some
258 participants: the cyprinid (“madrilla”) *Parachondrostoma miegii*; the minnow
259 (“piscardo”) *Phoxinus* spp.; the Iberian barbel (“barbo”) *Luciobarbus bocagei*; and, the
260 Iberian chub (“cacho”) *Squalius carolitertii*. Only one person interviewed in the Raíces
261 River region reported a native species to be an exotic species (*Anguilla anguilla*) (Table
262 S1).

263 A total of 32 participants (22.86% of the total) correctly identified at least one
264 exotic species. Seven exotic species were identified: the American crayfish (“cangrejo
265 americano”) *Procambarus clarkii*; the pond slider (“tortuga de Florida”) *Trachemys*
266 *scripta*; the common carp (“carpa”) *Cyprinus carpio*; the northern pike (“lucio”) *Esox*
267 *lucius*; the goldfish (“carpín”) *Carassius auratus*; the brook trout (“salvelino”)
268 *Salvelinus fontinalis*; and, the rainbow trout (“trucha arco iris”) *Oncorhynchus mykiss*
269 (Table S1). *P. clarkii* was reported as an exotic species by 10, 2, 2 and 3 participants
270 from the Piles, Nalón, Negro and Raíces river regions, respectively. In contrast, *C.*
271 *carpio* was more frequently reported as an exotic species in the Nalón River (8
272 participants) than in the Negro River (1 participant) and the other two rivers (no
273 participants).

274 A few participants (1 in Nalón, 1 in Negro and 3 in Piles) considered the Asian
275 carp *C. carpio* as a native species, and two participants (1 in Nalón and 1 in Piles)
276 regarded the American rainbow trout *O. mykiss* as being native (Table S1). The most
277 frequently reported NIS was *P. clarkii*, which was cited in the four rivers by a total of
278 17 citizens, followed by *C. carpio* (14 participants in three rivers) and *T. scripta* (6
279 participants in two rivers). These three species were also the most cited in recent media
280 releases (Table S2).

281 Significant differences in the knowledge about river species were not found
282 between genders and age groups (data not shown) with the exception of the number of
283 incorrect exotics (I_E). The mistakes about exotic species (I_E) were significantly different
284 between age groups (Kruskal-Wallis $H_c=3.97$; 3 degrees of freedom (df); $P=0.046$), and
285 there were clearly fewer mistakes by younger participants ($I_E=0\pm 0$) than by older
286 participants ($I_E=0.06\pm 0.24$). The rest of the data were pooled and organized by river.
287 Knowledge on native species (C_N) was significantly different among rivers (Kruskal-
288 Wallis $H_c=41.87$; 3 df; $P=4.27\times 10^{-9}$) (Table 2), and the knowledge levels were clearly
289 lower in the Raíces River region ($C_N=0.86\pm 0.60$) than in the rest of the regions (Figure
290 2). The knowledge of exotic species (C_E) was significantly lower (Kruskal-Wallis
291 $H_c=14.13$; 3 df; $P=0.003$) in Negro and Raíces than in the Nalón and Piles river regions
292 (Figure 2). Mistakes about exotic and native species (I_E and I_N , respectively) were
293 generally lower than those about correct species assigned to these categories, and
294 significant differences among rivers were not found (Kruskal-Wallis of $H_c=0.62$; 3 df;
295 $P=0.892$ and $H_c=4.07$; 3 df; $P=0.254$, respectively). The knowledge index, K_i , was
296 significantly different among river regions (Kruskal-Wallis of $H_c=32.97$; 3 df;
297 $P=3.27\times 10^{-7}$), and the K_i was lower in the Raíces River region than in the other river
298 regions accordingly (Figure 2), as fewer native species were reported from the Raíces

299 River. Significant differences between genders were not found for any question
300 regarding perception/opinion about NIS (data not shown). The two-way ANOVA that
301 considered age and river as factors revealed significant differences between ages and
302 among rivers for Question A, which was about the potential of NIS to adapt in the rivers
303 of the region (F=4.33 with P=0.039 for age; F= 2.86 with P=0.039 for river; F= 0.317
304 with P=0.813 for interaction).

305 Participants in the Nalón area and younger participants perceived, on average, a
306 higher capacity for the adaptation of NIS than other participant groups (Figure 3). For
307 Question D (demanding NIS eradication from Asturias rivers), highly significant
308 differences among rivers were found (F=0.034 with P=0.853 for age; F=8.922 with
309 $P=2 \times 10^{-5}$ for rivers; F=0.287 with P=0.835 for interaction). Participants interviewed in
310 the Raíces River region were less supportive of the eradication of exotic species than
311 those interviewed in other river regions (Figure 3). For the other two questions on how
312 much exotics affect native species (Question B) and how intense the changes perceived
313 in the river ecosystem are (Question C), significant differences were not found among
314 rivers nor between ages (data not shown).

315 Table 3 presents pairwise Spearman's r_s correlations in the dataset. The
316 knowledge index, K_i , was positively correlated with the knowledge about native and
317 exotic species (Table 3), as expected. Interestingly, after Bonferroni correction, the
318 knowledge index was positively correlated with the demand for the control actions
319 against exotic species, Question D ($P=3.21 \times 10^{-4}$). The number of correctly identified
320 native species (C_N) was also positively correlated with Question D (control of exotic
321 species) ($P=3.08 \times 10^{-5}$). The number of correctly identified exotic species (C_E) was
322 consistently positively correlated with the perception of the adaptation ability of the
323 exotic species, which was Question A ($P= 2.55 \times 10^{-3}$). As also expected, Question B
324 (opinion on how harmful NIS are to native species) was highly positively correlated
325 with Question D, which was the demand for NIS control ($P= 5.71 \times 10^{-8}$). The main
326 changes detected in the ecosystem by participants (Figure 4) were changes in the river
327 environment (59 participants). More participants from the Nalón River region (12) than
328 from the other zones detected changes in water quality; 7 of them reported improved
329 water quality. For river fauna, in the Negro River region, 11 participants noticed a
330 decrease in the *S. trutta* population in the region. In the river environment category,
331 more citizens in the Piles River region detected changes in the ecosystem, while in the
332 river infrastructure category, many Raíces River participants (11) reported a new
333 promenade near the riverbank.

334 Regarding the value of public knowledge used for early alerts of exotic species
335 in river systems, in this case study, *Acipenser sturio*, *Esox lucius* and *Luciobarbus*
336 *bocagei* were listed by different participants as occurring in the region although they
337 have not yet been found in biodiversity surveys in Asturias rivers (Ministerio de Medio
338 Ambiente 2007). On the other hand, the electrofishing survey detected the pond slider
339 *Trachemys scripta* (Table S1) in the river where the participants reported it. The species
340 is cataloged in the official list of exotic species, but until now it has been reported from
341 only isolated artificial ponds in Gijón and La Granda (Pleguezuelos 2002). Thus, this is
342 the first time the exotic pond slider was found in the wild in this region.

343 From the electrofishing survey, a total of 8 NIS were found in the region:
344 *Chondrostoma duriense*; *Cobitis paludica*; *Gobio lozanoi*, *Phoxinus* spp.; *Squalius*
345 *carolitertii*; *Carassius auratus*; *Procambarus clarkii*; and, *Trachemys scripta*. Seven
346 native species were sampled: *Anguilla anguilla*; *Chelon labrosus*; *Dicentrarchus*
347 *labrax*; *Mugil cephalus*; *Petromyzon marinus*; *Platichthys flesus*; and, *Salmo trutta*
348 (Table S1). The brown trout *Salmo trutta* was the only species found in all four rivers.

349 The Nalón River contained more NIS (five species), and the Raíces River exhibited the
350 highest proportion of NIS (three NIS out of a total of four species, 75%) (Table 4).

351 Comparing the aquatic fauna found from the electrofishing survey with the
352 knowledge of the local citizens (Table 4) revealed that the percentage of native species
353 recognized by locals in the four river regions was higher than the percentage of NIS. In
354 the Negro River region, where no NIS and only two native species were found from
355 electrofishing (Table 1S), participants recognized all species surveyed. In the Piles
356 River region, citizens were able to recognize 80% of the surveyed native species. In the
357 Raíces River region, participants recognized the native species but only one of three
358 exotic species. In the Nalón River region, citizens recognized 50% of the native species
359 sampled from the river (the same percentage of the official records).

360

361 **4. Discussion**

362

363 This case study illustrates the importance of considering the knowledge of citizens and
364 their opinions on treating biodiversity issues. Despite the relatively limited knowledge
365 about NIS, citizens were generally aware of their potential risks. Although only 22.9%
366 of the participants correctly recognized any NIS, as many as 73.6% were of the opinion
367 it is necessary to act against exotic species, and 67.9% believed that NIS could affect
368 native species. Accordingly, there was a positive attitude towards the eradication of the
369 NIS that affect native aquatic fauna. The results were similar to those found in Scotland
370 where 87% of the respondents supported the control and eradication of invasive species
371 (Bremner & Park, 2007).

372 In our particular case, the correspondence between real data, government reports
373 and citizen data was not accurate in relation to NIS; but, citizens recognized more than
374 50% of the native species surveyed by electrofishing in each river (Table 4), and they
375 were able to detect one exotic species in running waters in the wild, *Trachemys scripta*,
376 which was previously believed to only occur in artificial ponds (Pleguezuelos 2002)
377 (Table S1). The occurrence of species in the Raíces River was confirmed by
378 electrofishing in our study. This is a case where citizens reported a NIS that was
379 overlooked in official reports, and, as emphasized by many other authors working with
380 invasive species (Gallo & Waitt, 2011; Zenetos et al., 2013; Hawthorne et al., 2015;
381 Kobori et al., 2016; Maistrello et al., 2016), this result reinforces the importance of
382 counting on citizen scientists.

383 The local knowledge about native species was much greater than that about NIS.
384 More than 80% of the participants listed brown trout as a native species. Brown trout is
385 actually the dominant freshwater species in the region (e.g., Lobón-Cerviá, 2009) and
386 was the only species found from all four rivers considered in this study (Table S1).
387 Interestingly, such knowledge about the native fauna was highly and positively
388 correlated with the demand for NIS eradication (Table 3). Positive correlations between
389 local knowledge and awareness about biodiversity have been found by other authors in
390 Scotland, Chile and the Pyrenees (Bremner & Park, 2007; Loyau & Schmeller, 2017;
391 Zorondo-Rodríguez, Reyes-García, & Simonetti, 2014), and the results of this study are
392 along the same lines.

393 In the Raíces River region, the knowledge of the native and exotic species was
394 significantly lower than in the rest of river regions, as was the support of actions against
395 NIS. This could be explained by the lower quality environmental conditions in this
396 river. The Raíces River is a small narrow coastal stream (<2 meters wide), with very
397 reduced water flow. The local people may believe that there is not aquatic fauna in the
398 river and conservation efforts are not worthy there. The electrofishing survey revealed a

399 population of the native species *S. trutta*. It also revealed that the Raíces River is
400 invaded by NIS, since 75% of the species surveyed were exotics, including *Phoxinus*
401 spp., *P. clarkii* and *T. scripta*.

402 Better environmental education will improve the public awareness of NIS and
403 reduce the intentional release of some aquatic species (Zenetos et al., 2013). For
404 instance, *Carassius auratus*, which was found in Gijón, can be purchased in any pet
405 shop and is likely one of the cases of releases from pet owners (Elvira & Almodóvar,
406 2001; Maceda-Veiga, Domínguez-Domínguez, Escribano-Alacid, & Lyons, 2016), as
407 reported in the Pacific Northwest (Strecker, Campbell, & Olden, 2011), Iberian
408 Peninsula (Maceda-Veiga, Escribano-Alacid, de Sostoa, & García-Berthou, 2013), and
409 Czech Republic (Lusková, Lusk, Halačka, & Vetešník, 2010). The importance of good
410 environmental education is undeniable. Jordan, Gray, Howe, Brooks, and Ehrenfeld
411 (2011) showed a substantial change in behavior regarding invasive plants after citizens
412 acquired new knowledge about them. Environmental education would reduce the
413 misclassification of species and likely increase the reports of non-native species. In our
414 study, the species officially cataloged as exotics in Spain were considered NIS by the
415 participants, except for two fishes that were misidentified as native species by some
416 respondents: *Cyprinus carpio*; and, *Oncorhynchus mykiss*. These species are old
417 introductions since *C. carpio* was introduced to Spain in the 17th century (Elvira &
418 Almodóvar, 2001) and *O. mykiss* has been farmed in the region for more than 50 years
419 (Stanković, Crivelli, Snoj, Stankovi, & Snoj, 2015). People tend to be more aware of
420 recent introductions (García-Llorente et al., 2008). The most cited exotic in our study
421 was the American crayfish *P. clarkii*, which was identified in all four river regions. In
422 Gijón (Piles River), 10 participants out of the 35 identified *P. clarkii* as an invasive
423 species, compared with two participants in the Nalón and Negro river regions and three
424 in the Raíces River region. This is consistent with the higher awareness about recent
425 introductions (García-Llorente et al., 2008) because this species was found in an
426 artificial pond in downtown Gijón in June 2016, and the discovery was highly
427 publicized in the local newspapers (Table S2).

428 Although few people recognized any alien species in this study, 77.9% of the
429 participants were able to notice changes in the river ecosystem. Increasing the local
430 knowledge could help control non-native species. In general, citizen science programs
431 are cheaper and more affordable than research programs where scientists obtain the data
432 (Delaney et al., 2008). In our case, the cost would make the monitoring of the rivers and
433 the aquatic fauna at every moment throughout the region impossible without the help of
434 citizen science. As an example, in the Netherlands, Nunes and Van den Bergh (2004)
435 calculated that the benefits of a marine protection program far exceeded the costs with
436 the help of citizen science. Therefore, citizen science programs will help make research
437 cheaper and profitable, especially in this era when mobile phones and applications are
438 continuously renewed throughout the world (Newman et al., 2012). A strategy to
439 develop better responses to invasive species is publicly sharing the information
440 collected (Simpson et al., 2009), as is the case for the open-source atlas of invasive
441 plants of New England created in 2001, which offers presence/absence data and
442 contributes to many studies (Bois et al., 2011). However, it is necessary to create good
443 cyber infrastructure to manage the vast amounts of data from the citizen science
444 programs (Dickinson et al., 2012; Kobori et al., 2016).

445 On the other hand, molecular methods, such as environmental DNA (eDNA),
446 have been recently developed for the early detection of exotic species (Clusa et al.,
447 2016; Ficetola, Miaud, Pompanon, & Taberlet, 2008; Thomsen & Willerslev, 2015).
448 Together, eDNA and citizen science could be a promising tool to monitor and avoid the

449 spread of non-native species. For example, researchers trained 20 volunteers to
450 differentiate between the invasive pygmy mussel (*Xenostrobus securis*) and native
451 mussels in Asturias. In one day, volunteers were able to clean the affected area. The
452 eDNA tool made it possible to monitor the population in the region after the cleaning,
453 and the results demonstrated the success of the eradication process (Miralles, Dopico,
454 Devlo-Delva, & Garcia-Vazquez, 2016). Also, in the United Kingdom, the use of
455 volunteers to collect eDNA samples across the country helped to monitor the status of
456 the crested newt (*Triturus cristatus*) (Biggs et al., 2015).

457 Finally, efforts should focus on explaining the problems caused by NIS to the
458 local population and collaborating with the media to quickly divulge this knowledge to
459 the citizens, both of which could help detect new alien species that may come to the
460 region. Sharing research results with managers will help provide a better understanding
461 of the real fauna and the potential invaders, allowing for the design of better
462 management programs. In addition, the involvement of the general public through
463 citizen science, perhaps coupled with eDNA surveys, would be very helpful to prevent
464 the future spread of present and upcoming NIS in the region.

465 466 **5. Conclusion**

467
468 In this work, we detected how local citizens could be the first to detect significant
469 changes in the ecological environment of rivers and the introduction of any exotic
470 species. Early alert networks will contribute to transferring knowledge of any changes
471 detected to researchers and authorities for a rapid response. There is evidence that action
472 by citizens at an ecosystem level could keep the presence of non-native species under
473 control. For this reason, developing citizen science programs will increase public
474 interest in NIS intervention and may keep citizens in contact with scientific knowledge.
475 With better education about NIS, intentional releases may decrease, and people will be
476 more vigilant about their environment. Moreover, taking advantage of their enthusiasm
477 and motivation to participate in scientific research is also a strong incentive to share
478 scientific knowledge. In the case of Asturias, the local knowledge about non-indigenous
479 species is not accurate; but, the attitude towards these species makes the region a
480 promising candidate for focused education efforts to help preserve the fauna
481 biodiversity and protect against exotic species.

482 483 **6. Acknowledgements**

484
485 We thank Jorge Fernández De la Hoz for helping with the social survey. Laura Clusa
486 holds a PCTI Grant from the Asturias Regional Government, referenced BP14-145.
487 This work was supported by the Spanish project MINECO-13-CGL2013-42415-R and
488 the Asturias Regional Grant GRUPIN-2014-093. The EU RIA 689682 –AMBER
489 partially contributed to support this work. The authors do not have any conflict of
490 interest to declare. We are grateful to the two anonymous Reviewers who kindly helped
491 to improve this manuscript.

492 493 **7. Data Accessibility**

- 494
495 - Questionnaire is available in "Supplementary material_1".
496 - Data obtained from the 140 surveys are available in "Supplementary material_2".

497 - List of aquatic species and their status in Asturias together with the results from the
498 official inventory, electrofishing results and social survey in the 4 rivers are available in
499 "Supplementary material_3_Table S1"
500 - Compilation of data from official inventory, electrofishing results, social survey and
501 releases in public media regarding non indigenous species is available in
502 "Supplementary material_4_Table S2".
503 - All the supplementary material is available in the online repository figshare:
504 <https://figshare.com/s/cb7524b36edc574de412> doi: [10.6084/m9.figshare.5357671](https://doi.org/10.6084/m9.figshare.5357671)
505

506 8. References

- 507
- 508 Azzurro, E., Broglio, E., Maynou, F., & Bariche, M. (2013). Citizen science detects the
509 undetected: the case of *Abudefduf saxatilis* from the Mediterranean Sea.
510 *Management of Biological Invasions*, 4(2), 167–170.
511 <http://doi.org/http://dx.doi.org/10.3391/mbi.2013.4.2.10>
- 512 Biggs, J., Ewald, N., Valentini, A., Gaboriaud, C., Dejean, T., Griffiths, R. A., ... Dunn,
513 F. (2015). Using eDNA to develop a national citizen science-based monitoring
514 programme for the great crested newt (*Triturus cristatus*). *Biological Conservation*,
515 183, 19–28. <http://doi.org/10.1016/j.biocon.2014.11.029>
- 516 Bois, S. T., Silander, J. A., & Mehrhoff, L. J. (2011). Invasive Plant Atlas of New
517 England: The Role of Citizens in the Science of Invasive Alien Species Detection.
518 *BioScience*, 61(10), 763–770. <http://doi.org/10.1525/bio.2011.61.10.6>
- 519 Bremner, A., & Park, K. (2007). Public attitudes to the management of invasive non-
520 native species in Scotland. *Biological Conservation*, 139(3-4), 306–314.
521 <http://doi.org/10.1016/j.biocon.2007.07.005>
- 522 Carballo-Cárdenas, E. C., & Tobi, H. (2016). Citizen science regarding invasive lionfish
523 in Dutch Caribbean MPAs: Drivers and barriers to participation. *Ocean and*
524 *Coastal Management*, 133, 114–127.
525 <http://doi.org/10.1016/j.ocecoaman.2016.09.014>
- 526 Chown, S. L., Hodgins, K. A., Griffin, P. C., Oakeshott, J. G., Byrne, M., & Hoffmann,
527 A. A. (2015). Biological invasions, climate change and genomics. *Evolutionary*
528 *Applications*, 8(1), 23–46. <http://doi.org/10.1111/eva.12234>
- 529 Clusa, L., Ardura, A., Gower, F., Miralles, L., Tsartsianidou, V., Zaiko, A., & Garcia-
530 Vazquez, E. (2016). An Easy Phylogenetically Informative Method to Trace the
531 Globally Invasive *Potamopyrgus* Mud Snail from River's eDNA. *Plos One*,
532 11(10), e0162899. <http://doi.org/10.1371/journal.pone.0162899>
- 533 Crall, A. W., Newman, G. J., Jarnevich, C. S., Stohlgren, T. J., Waller, D. M., &
534 Graham, J. (2010). Improving and integrating data on invasive species collected by
535 citizen scientists. *Biological Invasions*, 12(10), 3419–3428.
536 <http://doi.org/10.1007/s10530-010-9740-9>
- 537 Crall, A. W., Newman, G. J., Stohlgren, T. J., Holfelder, K. A., Graham, J., & Waller,
538 D. M. (2011). Assessing citizen science data quality: An invasive species case
539 study. *Conservation Letters*, 4(6), 433–442. <http://doi.org/10.1111/j.1755-263X.2011.00196.x>
- 540
- 541 Delaney, D. G., Sperling, C. D., Adams, C. S., & Leung, B. (2008). Marine invasive
542 species: Validation of citizen science and implications for national monitoring
543 networks. *Biological Invasions*, 10(1), 117–128. <http://doi.org/10.1007/s10530-007-9114-0>
544

- 545 De la Hoz, J. (2006). La gestión de la pesca continental en Asturias (España). Capturas,
546 repoblaciones y actuaciones de mejora del medio. Demanda de cotos y licencias.
547 *Bol. Cien. Nat. RIDEA*, Oviedo.
- 548 Devictor, V., Whittaker, R. J., & Beltrame, C. (2010). Beyond scarcity: Citizen science
549 programmes as useful tools for conservation biogeography. *Diversity and*
550 *Distributions*, 16(3), 354–362. <http://doi.org/10.1111/j.1472-4642.2009.00615.x>
- 551 Dickinson, J. L., Shirk, J., Bonter, D., Bonney, R., Crain, R. L., Martin, J., ... Purcell,
552 K. (2012). The current state of citizen science as a tool for ecological research and
553 public engagement. *Frontiers in Ecology and the Environment*, 10(6), 291–297.
554 <http://doi.org/10.1890/110236>
- 555 Dickinson, J., Zuckerberg, B., & Bonter, D. (2010). Citizen Science as an Ecological
556 Research Tool: Challenges and Benefits. *Annual Review of Ecology, Evolution and*
557 *Systematics*, 41(1), 149–172. [http://doi.org/10.1146/annurev-ecolsys-102209-](http://doi.org/10.1146/annurev-ecolsys-102209-144636)
558 [144636](http://doi.org/10.1146/annurev-ecolsys-102209-144636)
- 559 Elvira, B., & Almodóvar, A. (2001). Freshwater fish introductions in Spain : facts and
560 figures at the beginning of the 21st century. *Journal of Fish Biology*,
561 59(Supplement A), 323–331. <http://doi.org/10.1006/jfbi.2001.1753>
- 562 Falk, B. G., Snow, R. W., & Reed, R. N. (2016). Prospects and Limitations of Citizen
563 Science in Invasive Species Management: A Case Study with Burmese Pythons in
564 Everglades National Park. *Southeastern Naturalist*, 15(sp8), 89–102.
565 <http://doi.org/10.1656/058.015.sp806>
- 566 Ficetola, G. F., Miaud, C., Pompanon, F., & Taberlet, P. (2008). Species detection using
567 environmental DNA from water samples. *Biology Letters*, 4(4), 423–425.
568 <http://doi.org/10.1098/rsbl.2008.0118>
- 569 Gallo, T., & Waitt, D. (2011). Creating a Successful Citizen Science Model to Detect
570 and Report Invasive Species. *BioScience*, 61(6), 459–465.
571 <http://doi.org/10.1525/bio.2011.61.6.8>
- 572 García-Llorente, M., Martín-López, B., González, J. A., Alcorlo, P., & Montes, C.
573 (2008). Social Perceptions of the Impacts and Benefits of Invasive Alien Species :
574 Implications for Management. *Biological Conservation*, 141, 2969–2983.
575 <http://doi.org/10.1016/j.biocon.2008.09.003>
- 576 Hammer, Ø., Harper, D. A. T., & Ryan, P. D. (2001). PAST: Paleontological Statistics
577 Software Package for Education and Data Analysis. *Palaeontologia Electronica*,
578 4(1)(1), 1–9. <http://doi.org/10.1016/j.bcp.2008.05.025>
- 579 Havel, J. E., Kovalenko, K. E., Thomaz, S. M., Amalfitano, S., & Kats, L. B. (2015).
580 Aquatic invasive species: challenges for the future. *Hydrobiologia*, 750(1), 147–
581 170. <http://doi.org/10.1007/s10750-014-2166-0>
- 582 Hawthorne, T. L., Elmore, V., Strong, A., Bennett-Martin, P., Finnie, J., Parkman, J., ...
583 Reed, J. (2015). Mapping non-native invasive species and accessibility in an urban
584 forest: A case study of participatory mapping and citizen science in Atlanta,
585 Georgia. *Applied Geography*, 56, 187–198.
586 <http://doi.org/10.1016/j.apgeog.2014.10.005>
- 587 Hulme, P. E. (2009). Trade, transport and trouble: Managing invasive species pathways
588 in an era of globalization. *Journal of Applied Ecology*, 46(1), 10–18.
589 <http://doi.org/10.1111/j.1365-2664.2008.01600.x>
- 590 Ingwell, L. L., & Preisser, E. L. (2011). Using citizen science programs to identify host
591 resistance in pest-invaded forests. *Conservation Biology*, 25(1), 182–188.
592 <http://doi.org/10.1111/j.1523-1739.2010.01567.x>
- 593 Jordan, R. C., Gray, S. A., Howe, D. V., Brooks, W. R., & Ehrenfeld, J. G. (2011).
594 Knowledge Gain and Behavioral Change in Citizen-Science Programs.

595 *Conservation Biology*, 25(6), 1148–1154. <http://doi.org/10.1111/j.1523->
596 1739.2011.01745.x

597 Kobori, H., Dickinson, J. L., Washitani, I., Sakurai, R., Amano, T., Komatsu, N., ...
598 Miller-Rushing, A. J. (2016). Citizen science: a new approach to advance ecology,
599 education, and conservation. *Ecological Research*, 31(1), 1–19.
600 <http://doi.org/10.1007/s11284-015-1314-y>

601 Lobón-Cerviá, J. (2009). Why, when and how do fish populations decline, collapse and
602 recover? the example of brown trout (*Salmo trutta*) in Rio Chaballos (northwestern
603 Spain). *Freshwater Biology*, 54(6), 1149–1162. <http://doi.org/10.1111/j.1365->
604 2427.2008.02159.x

605 Loyau, A., & Schmeller, D. S. (2017). Positive sentiment and knowledge increase
606 tolerance towards conservation actions. *Biodiversity and Conservation*, 26(2), 461–
607 478. <http://doi.org/10.1007/s10531-016-1253-0>

608 Lusková, V., Lusk, S., Halačka, K., & Vetešník, L. (2010). *Carassius auratus gibelio*-
609 The Most Successful Invasive Fish in Waters of the Czech Republic. *Russian*
610 *Journal of Biological Invasions*, 1(3), 176–180.
611 <http://doi.org/10.1134/S2075111710030069>

612 Maceda-Veiga, A., Domínguez-Domínguez, O., Escribano-Alacid, J., & Lyons, J.
613 (2016). The aquarium hobby: can sinners become saints in freshwater fish
614 conservation? *Fish and Fisheries*, 17(3), 860–874. <http://doi.org/10.1111/faf.12097>

615 Maceda-Veiga, A., Escribano-Alacid, J., de Sostoa, A., & García-Berthou, E. (2013).
616 The aquarium trade as a potential source of fish introductions in southwestern
617 Europe. *Biological Invasions*, 15(12), 2707–2716. <http://doi.org/10.1007/s10530->
618 013-0485-0

619 Maistrello, L., Dioli, P., Bariselli, M., Mazzoli, G. L., & Giacalone-Forini, I. (2016).
620 Citizen science and early detection of invasive species: phenology of first
621 occurrences of *Halyomorpha halys* in Southern Europe. *Biological Invasions*,
622 18(11), 3109–3116. <http://doi.org/10.1007/s10530-016-1217-z>

623 Ministerio de Medio Ambiente. (2007). La invasión de Especies exóticas en los ríos.
624 Madrid, 124p.

625 Miralles, L., Dopico, E., Devlo-Delva, F., & Garcia-Vazquez, E. (2016). Controlling
626 populations of invasive pygmy mussel (*Xenostrobus securis*) through citizen
627 science and environmental DNA. *Marine Pollution Bulletin*, 110(1), 127–132.
628 <http://doi.org/10.1016/j.marpolbul.2016.06.072>

629 Newman, G., Wiggins, A., Crall, A., Graham, E., Newman, S., & Crowston, K. (2012).
630 The future of Citizen science: Emerging technologies and shifting paradigms.
631 *Frontiers in Ecology and the Environment*, 10(6), 298–304.
632 <http://doi.org/10.1890/110294>

633 Nunes, P. A., & Van den Bergh, J. C. (2004). Can people value protection against
634 invasive marine species? Evidence from a joint TC-CV survey in the Netherlands.
635 *Environmental and Resource Economics*, 28, 517–532. <http://doi.org/10.1023/B>
636 Occhipinti-Ambrogi, A., Galil, B. S. (2004). A uniform terminology on bioinvasions: A
637 chimera or an operative tool? *Marine Pollution Bulletin*, 49(9-10), 688–94.

638 Pleguezuelos, J. M. (2002). Las especies introducidas de anfibios y reptiles. Pp: 503-
639 532. En: Pleguezuelos, J.M., Márquez, R., Lizana, M. (Eds.). *Atlas y Libro Rojo de*
640 *los Anfibios y Reptiles de España*. Dirección General de Conservación de la
641 Naturaleza-Asociación Herpetológica Española (2ª impresión). Madrid

642 Silvertown, J. (2009). A new dawn for citizen science. *Trends in Ecology & Evolution*,
643 24(9), 467–71. <http://doi.org/10.1016/j.tree.2009.03.017>

644 Simpson, A., Jarnevich, C., Madsen, J., Westbrooks, R., Fournier, C., Mehrhoff, L., ...
645 Sellers, E. (2009). Invasive species information networks: collaboration at multiple
646 scales for prevention, early detection, and rapid response to invasive alien species.
647 *Biodiversity*, 10(2-3), 5–13. <http://doi.org/10.1080/14888386.2009.9712839>

648 Stanković, D., Crivelli, A. J., Snoj, A., Stankovi, D., & Snoj, A. S. (2015). Rainbow
649 trout in Europe: Introduction, naturalization, and impacts. *Reviews in Fisheries*
650 *Science & Aquaculture*, 23(1), 39–71.
651 <http://doi.org/10.1080/23308249.2015.1024825>

652 Strecker, A. L., Campbell, P. M., & Olden, J. D. (2011). The aquarium trade as an
653 invasion pathway in the Pacific Northwest. *Fisheries*, 36(2), 74–85.
654 <http://doi.org/10.1577/03632415.2011.10389070>

655 Sullivan, B. L., Aycrigg, J. L., Barry, J. H., Bonney, R. E., Bruns, N., Cooper, C. B., ...
656 Kelling, S. (2014). The eBird enterprise: An integrated approach to development
657 and application of citizen science. *Biological Conservation*, 169, 31–40.
658 <http://doi.org/10.1016/j.biocon.2013.11.003>

659 Thomsen, P. F., & Willerslev, E. (2015). Environmental DNA - An emerging tool in
660 conservation for monitoring past and present biodiversity. *Biological*
661 *Conservation*, 183, 4–18. <http://doi.org/10.1016/j.biocon.2014.11.019>

662 Tulloch, A. I. T., Possingham, H. P., Joseph, L. N., Szabo, J., & Martin, T. G. (2013).
663 Realising the full potential of citizen science monitoring programs. *Biological*
664 *Conservation*, 165, 128–138. <http://doi.org/10.1016/j.biocon.2013.05.025>

665 Zenetos, A., Koutsogiannopoulos, D., Ovalis, P., & Poursanidis, D. (2013). The role
666 played by citizen scientists in monitoring marine alien species in Greece. *Cahiers*
667 *de Biologie Marine*, 54(3), 419–426.

668 Zorondo-Rodríguez, F., Reyes-García, V., & Simonetti, J. A. (2014). Conservation of
669 biodiversity in private lands : are Chilean landowners willing to keep threatened
670 species in their lands? *Revista Chilena de Historia Natural*, 1, 4.
671 <http://doi.org/10.1186/0717-6317-1-4>

672

674

675 9. Tables

676

677

678 Table 1. **Sample for social survey.** Number of citizens classified by gender and age in
679 each river is shown.

680

681

	Nalón		Negro		Piles		Raíces		Total by age
	Men	Women	Men	Women	Men	Women	Men	Women	
20-30	0	2	0	0	3	2	0	2	9
30-40	5	5	2	2	1	2	3	3	23
40-50	6	3	4	2	2	4	3	5	29
50-60	5	1	4	4	5	3	6	3	31
>60	8	0	11	6	7	6	7	3	48
Total by river	35		35		35		35		140

682

683

684

685
 686
 687
 688
 689
 690
 691
 692
 693

Table 2. Kruskal-Wallis tests of the results analyzed by basin, age or gender. Results are based on averages across the different sample groups. Significant *P* values for differences among or between sample medians are in bold. C_N and C_E are correct native and correct exotic species identified. I_N and I_E are incorrect native and incorrect exotic species identified.

	C _N	I _N	C _E	I _E	Knowledge index (Ki)	A- Exotic adaptation	B- Harm to natives	C- Ecosystem changes	D-Exotics's removal
Basin	<0.001	0.254	0.003	0.892	<0.001	0.041	0.238	0.061	<0.001
Age (<50 and >50)	0.625	0.942	0.262	0.046	0.660	0.041	0.557	0.308	0.784
Gender	0.490	0.283	0.297	0.947	0.385	0.310	0.269	0.634	0.780

694
 695
 696

697 Table 3. **Spearman's rs correlation results of the 140 surveys.** The rs and p values are
 698 below and above the diagonal, respectively. Significant correlations (after Bonferroni
 699 correction) are indicated in bold and significant p-values are highlighted in grey. C_N and
 700 C_E are correct native and correct exotic species identified. I_N and I_E are incorrect native
 701 and incorrect exotic species identified. Ki = knowledge index. A-Exotic adaptation is
 702 the answer to the ability of exotic species to adapt in Asturian rivers; B- harm to natives
 703 is the answer to the ability of exotic species to affect native fauna; C- Ecosystem
 704 changes is the answer to detection of changes in the ecosystem by the citizens and D-
 705 Exotic's removal is the answer to the necessity of taking action against exotic species.
 706
 707

	C _N	I _N	C _E	I _E	A- Exotic adaptation	B- Harm to natives	C- Ecosystem changes	D- Exotics's removal	Knowledge index (Ki)
C _N		0.439	0.150	0.143	0.466	0.083	0.032	3.08x10 ⁻⁵	1.42x10 ⁻³⁷
I _N	0.066		0.012	0.452	0.468	0.116	0.442	0.084	0.023
C _E	0.122	0.213		0.391	2.55x10 ⁻³	0.149	0.343	0.212	2.14x10 ⁻⁸
I _E	0.125	-0.064	0.073		0.223	0.675	0.228	0.724	0.857
A- Exotic adaptation	-0.062	-0.062	0.253	0.104		0.022	0.715	0.021	0.347
B- Harm to natives	0.147	0.134	0.123	0.036	0.193		0.959	5.71x10 ⁻⁸	0.087
C- Ecosystem changes	0.182	0.065	0.081	0.103	0.031	0.004		0.243	0.135
D- Exotics's removal	0.344	0.146	0.106	0.030	0.194	0.439	0.099		3.21x10 ⁻⁴
Knowledge index (Ki)	0.835	-0.192	0.452	-0.015	0.080	0.145	0.127	0.300	

708
 709
 710

711 Table 4. **Comparison between real aquatic fauna, official records and local citizens'**
712 **data.** Two results are considered per river, the number of correct species (native, exotic
713 and total species) listed by volunteers over the total number of species in the region
714 based on official records (over region), and the number of correct species listed by
715 volunteers over the number of species found in the electrofishing survey (over survey).
716 In parenthesis percentage of species recognized by locals over region and over survey is
717 shown.
718

	Region (N)	Nalón			Negro			Piles			Raíces		
		Electrofishing Survey	Recognized by locals		Electrofishing Survey	Recognized by locals		Electrofishing Survey	Recognized by locals		Electrofishing Survey	Recognized by locals	
			Over region	Over survey		Over region	Over survey		Over region	Over survey		Over region	Over survey
Native	12	4	6 (50%)	2 (50%)	2	5 (41.7%)	2 (100%)	5	6 (50%)	4 (80%)	1	4 (33.3%)	1 (100%)
Exotic	16	5	6 (37.5%)	2 (40%)	0	6 (37.5%)	0 (0%)	2	4 (25%)	0 (0%)	3	2 (12.5%)	1 (33.3%)
Total	28	9	12 (42.9%)	4 (44.4%)	2	11 (39.3%)	2 (100%)	7	10 (35.7%)	4 (57.1%)	4	6 (21.4%)	2 (50%)

719

720

721

722 10. Figures

723

724 **Figure 1. Map showing the sampling sites of the four rivers: Nalón, Negro, Piles and**
725 **Raíces**

726

727

728

729

730

731

732

733

734

735

736

737

738

739

740

741

742

743

744

745

746

747

748

749

750

751

752

753

754

755


756

757

758


759

760




761 Figure 2. **Mean and standard deviation of public knowledge per basin.** Correct
 762 natives (C_N), incorrect natives (I_N), correct exotics (C_E), incorrect exotics (I_E) and
 763 knowledge index (K_i) are shown.

764
 765
 766
 767
 768
 769
 770
 771
 772
 773
 774
 775
 776
 777
 778
 779
 780
 781
 782



783 **Figure 3. Mean and standard deviation of perception issues about exotic species per**
 784 **basin and age.** A-"Exotic adaptation" is the answer to the ability of exotic species to
 785 adapt in Asturian rivers; B- "harm to natives" is the answer to the ability of exotic
 786 species to affect native fauna; C- "Ecosystem changes" is the answer to detection of
 787 changes in the ecosystem by the citizens and D- "Exotic's removal" is the answer to the
 788 necessity of taking action against exotic species.



799
 800
 801
 802
 803
 804
 805
 806
 807
 808
 809
 810

811 Figure 4. **Changes in the environment reported by the citizens in the four rivers**
 812 **surveyed.** Four groups of changes were considered: changes water quality (including
 813 water flow, algae, or sediments), changes regarding river fauna (less *S. trutta*, more
 814 *Mugil cephalus*); infrastructure (new ponds, dams, promenade); and environment
 815 (cleaner environment, more pollution, more vegetation). Number of citizens in each
 816 place expressing each kind of change is shown.

817
 818
 819

